

# Target Corporation

**-PE Product Labels –**

**-continued**





See new warnings information & directions  
Compare to active ingredients in  
Theraflu® Nighttime Severe Cold & Cough\*

NDC 11673-113-07

# nighttime severe cold, cough and flu

**acetaminophen** (pain reliever/fever reducer)  
**diphenhydramine HCl** (antihistamine/cough suppressant)  
**phenylephrine HCl** (nasal decongestant)

nasal congestion, cough,  
runny nose, sneezing,  
body ache, sore throat pain,  
headache, fever

honey lemon infused  
with chamomile and  
white tea flavors



6 PACKETS



# nighttime severe cold, cough and flu acetaminophen

(pain reliever/fever reducer)  
**diphenhydramine HCl**  
(antihistamine/cough suppressant)  
**phenylephrine HCl**  
(nasal decongestant)



\*This product is not manufactured or  
distributed by Novartis Consumer Health, Inc.  
or their affiliates, owner of the registered  
trademark Theraflu®

094 03 0598 R00 ID305413  
Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in Israel  
©2015 Target Brands, Inc.  
Shop Target.com





READ ALL WARNINGS AND DIRECTIONS ON CARTON BEFORE USE.  
KEEP CARTON FOR REFERENCE. DO NOT DISCARD.

## Drug Facts

### Active ingredients (in each packet)

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine hydrochloride 25 mg.....	Antihistamine/cough suppressant
Phenylephrine hydrochloride 10 mg.....	Nasal decongestant

### Purposes

### Uses

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - minor sore throat pain
  - headache
  - nasal and sinus congestion
  - runny nose
  - sneezing
  - itchy nose or throat
  - itchy, watery eyes due to hay fever
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

### Warnings

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- in a child under 4 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- a sodium-restricted diet
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema, asthma or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

### Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

### When using this product

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

TAMPER EVIDENT INNER PACKET.  
DO NOT USE IF SEALED PACKET IS TORN OR BROKEN

## Drug Facts (continued)

### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- symptoms do not get better or worsen
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health care professional before use.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- do not use more than directed
- take every 4 hours; do not take more than 5 packets in 24 hours unless directed by a doctor

Age	Dose
<b>children under 4 years of age</b>	<b>do not use</b>
children 4 to under 12 years of age	do not use unless directed by a doctor
adults and children 12 years of age and over	one packet

- dissolve contents of one packet into 8 oz. hot water; sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water; stir briskly before and after heating. Do not overheat.

### Other information




- each packet contains:** potassium 4 mg, sodium 27 mg
- phenylketonurics:** contains phenylalanine 34 mg per packet
- store at controlled room temperature 20°-25°C (68°-77°F). Protect product from heat and moisture.

### Inactive ingredients

acesulfame potassium, aspartame, citric acid anhydrous, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, flavors, maltodextrin, sodium citrate anhydrous, sucrose, and pregelatinized starch.

### Questions or Comments?

Call 1-800-910-6874





NDC 11673-789-40

# nighttime severe cold and flu honey flavor

**MAX**  
STRENGTH**acetaminophen**

(pain reliever/fever reducer)

phenylephrine HCl

(nasal decongestant)

dextromethorphan HBr

(cough suppressant)

doxylamine succinate

(antihistamine)

headache, fever, sore throat,

minor aches and pains

sneezing, runny nose

nasal/sinus congestion

and sinus pressure

cough



12 FL OZ (355 mL)

: 86S40 UW F1

**PARENTS:**

Learn about teen medicine abuse

www.StopMedicineAbuse.org

**DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING**

094 14 0003 R00 C-001744-01-067

Dist. by Target Corp., Mpls, MN 55403

**Drug Facts****Active ingredients**  
(in each 15 mL)

Active ingredients	Purpose
Acetaminophen	Pain reliever/
325 mg.....	fever reducer
Dextromethorphan HBr	
10 mg.....	Cough suppressant
Doxylamine Succinate 6.25 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ cough to help you sleep ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ runny nose and sneezing ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage

**Warnings**

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**PEEL BACK AT  
CORNER FOR MORE  
INFORMATION**

: 86S40 UW B1

**Drug Facts**  
(continued)

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ to make a child sleepy

**Ask a doctor before use if you have** ■ liver disease ■ heart disease ■ high blood pressure ■ glaucoma ■ thyroid disease ■ diabetes ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating

**Drug Facts**  
(continued)

■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

**WESIVE AREA  
• NO VARNISH • NO TYPE**

**ADHESIVE AREA  
• NO VARNISH • NO TYPE**

8118

**AD**  
**NO COATING**

due to an enlarged prostate gland  
 ■ a breathing problem such as emphysema or chronic bronchitis  
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema.

Ask a doctor or pharmacist before use if you are  
 ■ taking the blood thinning drug warfarin  
 ■ taking sedatives or tranquilizers

When using this product  
 ■ do not use more than directed  
 ■ excitability may occur, especially in children  
 ■ marked drowsiness may occur

**NO COATING**

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

**Other information** ■ each 15 mL contains: sodium 15 mg ■ store at 20-25°C (68-77°F)

**Inactive ingredients** anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

**Questions? Call**  
1-888-547-7400

TM & ©2022  
Target Brands, Inc.

daytime  
**severe**  
cold and flu

Compare to active  
ingredients in Vicks®  
DayQuil® Severe Cold & Flu\*

daytime  
**severe**  
cold and flu

acetaminophen  
(pain reliever/fever reducer)  
dextromethorphan HBr  
(cough suppressant)  
guaifenesin (expectorant)  
phenylephrine HCl  
(nasal decongestant)

headache, fever, sore throat,  
minor aches and pains,  
nasal/sinus congestion  
and sinus pressure, cough,  
chest congestion  
non-drowsy  
alcohol free



12 FL OZ (355 mL)

nighttime  
**severe**  
cold and flu

Compare to active  
ingredients in Vicks®  
NyQuil® Severe Cold & Flu\*

value  
pack

nighttime  
**severe**  
cold and flu

acetaminophen  
(pain reliever/fever reducer)  
dextromethorphan HBr  
(cough suppressant)  
doxylamine succinate  
(antihistamine)  
phenylephrine HCl  
(nasal decongestant)

headache, fever, sore throat,  
minor aches and pains,  
nasal/sinus congestion and  
sinus pressure, sneezing,  
runny nose, cough  
alcohol free



12 FL OZ (355 mL)

MAX  
STRENGTH



MAX  
STRENGTH



nighttime  
**severe**  
cold and flu

DO NOT USE IF PRINTED NECKBAND IS  
BROKEN OR MISSING

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

094 14 0070 R00  
C-001744-01-093



100% satisfaction guaranteed or your money back.

Distributed by Target Corporation  
Minneapolis, MN 55403  
TM & ©2022 Target Brands, Inc.

\*These products are not manufactured or  
distributed by Procter & Gamble, distributor of  
Vicks® DayQuil® Severe Cold & Flu and Vicks®



3120

## Nighttime Severe Cold &amp; Flu

Drug Facts	
Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant
<b>Uses</b> temporarily relieves common cold/flu symptoms:	
<ul style="list-style-type: none"> <li>nasal congestion</li> <li>sinus congestion and pressure</li> <li>minor aches and pains</li> <li>headache</li> <li>fever</li> <li>sore throat</li> <li>runny nose and sneezing</li> <li>cough due to minor throat and bronchial irritation</li> <li>cough to help you sleep</li> <li>reduces swelling of nasal passages</li> <li>promotes nasal and/or sinus drainage</li> <li>temporarily restores freer breathing through the nose</li> </ul>	
<b>Warnings</b>	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:	
<ul style="list-style-type: none"> <li>more than 4,000 mg of acetaminophen in 24 hours</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks every day while using this product</li> </ul>	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
<ul style="list-style-type: none"> <li>skin redness</li> <li>hives</li> <li>rash</li> </ul>	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
<b>Do not use</b>	
<ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients</li> </ul>	
<b>Ask a doctor before use if you have:</b>	
<ul style="list-style-type: none"> <li>liver disease</li> <li>heart disease</li> <li>high blood pressure</li> <li>thyroid disease</li> <li>diabetes</li> <li>glaucoma</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>a breathing problem such as emphysema or chronic bronchitis</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>a sodium-restricted diet</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are:</b>	
<ul style="list-style-type: none"> <li>taking sedatives or tranquilizers</li> <li>taking the blood thinning drug warfarin</li> </ul>	
<b>When using this product:</b>	
<ul style="list-style-type: none"> <li>do not use more than directed</li> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>avoid alcoholic drinks</li> <li>be careful when driving a motor vehicle or operating machinery</li> <li>alcohol, sedatives, and tranquilizers may increase drowsiness</li> </ul>	
<b>Stop use and ask a doctor if:</b>	
<ul style="list-style-type: none"> <li>you get nervous, dizzy or sleepless</li> <li>pain, nasal congestion, or cough gets worse or lasts more than 7 days</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts.</li> </ul>	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>take only as directed - see Overdose warning</li> <li>only use the dose cup provided</li> <li>do not exceed 4 doses per 24 hrs</li> </ul>	
adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use
<b>Other information</b>	
<ul style="list-style-type: none"> <li>each 30 mL contains sodium 41 mg</li> <li>store at 20°-25°C (68°-77°F)</li> </ul>	
<b>Inactive ingredients</b> anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum	
<b>Questions?</b> Call 1-888-547-7400	

## Daytime Severe Cold &amp; Flu

Drug Facts	
Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant
<b>Uses</b> temporarily relieves common cold/flu symptoms:	
<ul style="list-style-type: none"> <li>nasal congestion</li> <li>sinus congestion and pressure</li> <li>cough due to minor throat and bronchial irritation</li> <li>minor aches and pains</li> <li>headache</li> <li>fever</li> <li>sore throat</li> <li>reduces swelling of nasal passages</li> <li>temporarily restores freer breathing through the nose</li> <li>promotes nasal and/or sinus drainage</li> <li>helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of both mucus and make coughs more productive</li> </ul>	
<b>Warnings</b>	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if:	
<ul style="list-style-type: none"> <li>adult takes more than 4,000 mg of acetaminophen in 24 hours</li> <li>child takes more than 5 doses in 24 hours</li> <li>taken with other drugs containing acetaminophen</li> <li>adult has 3 or more alcoholic drinks every day while using this product</li> </ul>	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
<ul style="list-style-type: none"> <li>skin redness</li> <li>hives</li> <li>rash</li> </ul>	
If a skin reaction occurs, stop use and seek medical help right away.	
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.	
<b>Do not use</b>	
<ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</li> <li>if you have ever had an allergic reaction to this product or any of its ingredients</li> </ul>	
<b>Ask a doctor before use if you have:</b>	
<ul style="list-style-type: none"> <li>liver disease</li> <li>heart disease</li> <li>thyroid disease</li> <li>diabetes</li> <li>high blood pressure</li> <li>trouble urinating due to an enlarged prostate gland</li> <li>cough that occurs with too much phlegm (mucus)</li> <li>persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</b>	
<b>When using this product:</b>	
<ul style="list-style-type: none"> <li>do not use more than directed</li> <li>excitability may occur, especially in children</li> <li>marked drowsiness may occur</li> <li>avoid alcoholic drinks</li> <li>be careful when driving a motor vehicle or operating machinery</li> <li>alcohol, sedatives, and tranquilizers may increase drowsiness</li> </ul>	
<b>Stop use and ask a doctor if:</b>	
<ul style="list-style-type: none"> <li>you get nervous, dizzy or sleepless</li> <li>pain, nasal congestion, or cough gets worse or lasts more than 7 days (children) or 7 days (adults)</li> <li>fever gets worse or lasts more than 3 days</li> <li>redness or swelling is present</li> <li>new symptoms occur</li> <li>cough comes back or occurs with rash or headache that lasts.</li> </ul>	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
<b>Directions</b>	
<ul style="list-style-type: none"> <li>take only as directed - see Overdose warning</li> <li>only use the dose cup provided</li> <li>do not exceed 4 doses per 24 hrs</li> </ul>	
adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use
<b>Other information</b>	
<ul style="list-style-type: none"> <li>each 15 mL contains sodium 6 mg</li> <li>store at 20°-25°C (68°-77°F). Do not refrigerate.</li> </ul>	
<b>Inactive ingredients</b> butylated hydroxyanisole, edetate disodium, FD&C yellow #6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum	
<b>Questions?</b> Call 1-888-547-7400	

daytime  
**severe**  
**cold and flu**

up&amp;up

**Drug Facts** (continued)

■ taking the blood thinning drug warfarin

**When using this product** ■ **do not use more than directed** ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed – see Overdose warning  
■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information** ■ each 30 mL contains: sodium 44 mg  
■ store at 20-25°C (68-77°F)

**Inactive ingredients** alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C yellow #6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

**Questions? Call 1-888-547-7400**

\*This product is not manufactured or distributed by Procter & Gamble, distributor of Vicks® NyQuil® Severe Cold & Flu.

**GLUTEN FREE**

094 14 0019 R02  
IDC-000021-01-193

Dist. by Target Corp.  
Mpls., MN 55403  
©2016 Target Brands, Inc.

**PARENTS:**

Learn about teen medicine abuse

www.StopMedicineAbuse.org

3

70030 62793

6

NDC 11673-012-40

Compare to active ingredients in  
Vicks® NyQuil® Severe Cold & Flu\*

## nighttime severe cold and flu

acetaminophen (pain reliever/fever reducer)  
dextromethorphan HBr (cough suppressant)  
doxylamine succinate (antihistamine)  
phenylephrine HCl (nasal decongestant)



headache, fever, sore throat,  
minor aches and pains  
nasal/sinus congestion  
and sinus pressure  
sneezing and runny nose  
cough

ALCOHOL 10%

12 FL OZ (355 mL)

**MAX  
STRENGTH**

**Drug Facts** **DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING****Active ingredients (in each 30 mL)**

Acetaminophen 650 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Doxylamine succinate 12.5 mg.....	Antihistamine
Phenylephrine HCl 10 mg.....	Nasal decongestant

**Purpose**

**Uses** temporarily relieves common cold/flu symptoms:

■ nasal congestion ■ sinus congestion and pressure ■ headache ■ fever  
■ minor aches and pains ■ sore throat ■ runny nose and sneezing  
■ cough due to minor throat and bronchial irritation ■ cough to help you sleep  
■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage  
■ temporarily restores freer breathing through the nose

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours  
■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have** ■ liver disease ■ heart disease  
■ high blood pressure ■ thyroid disease ■ a sodium-restricted diet  
■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma  
■ cough that occurs with too much phlegm (mucus)  
■ trouble urinating due to an enlarged prostate gland  
■ persistent or chronic cough as occurs with  
smoking, asthma, or emphysema ■ diabetes

**Ask a doctor or pharmacist before use if you are**  
■ taking sedatives or tranquilizers

: 82940 UW F2







3123

Compare to active ingredients in Theraflu® Severe Cold\*

## daytime multi-symptom severe cold

**acetaminophen**  
(pain reliever/fever reducer)

**dextromethorphan HBr**  
(cough suppressant)

**phenylephrine HCl**  
(nasal decongestant)

nasal congestion, cough, body ache, sore throat pain, headache, fever

serve hot

**green tea and honey lemon**  
flavors



**6**

PACKETS

6 MULTI-SYMPTOM PACKETS

NDC 11673-877-55

Compare to active ingredients in Theraflu® Nighttime Severe Cold & Cough\*

value pack

## nighttime severe cold and cough

**acetaminophen**  
(pain reliever/fever reducer)

**diphenhydramine HCl**  
(antihistamine/cough suppressant)

**phenylephrine HCl**  
(nasal decongestant)

nasal congestion, cough, runny nose, sneezing, body ache, sore throat pain, headache, fever

serve hot

**honey lemon**  
infused with white tea flavors



**6**

PACKETS

6 NIGHTTIME PACKETS

DO NOT TAKE BOTH PRODUCTS AT THE SAME TIME OR TAKE MORE THAN 5 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

DO NOT TAKE A DOSE OF THE NIGHTTIME PRODUCT SOONER THAN 4 HOURS AFTER THE LAST DOSE OF MULTI-SYMPTOM PRODUCT UNLESS DIRECTED BY YOUR DOCTOR.

Daytime Multi-Symptom Severe Cold

### Drug Facts

#### Active Ingredients (in each packet)

Acetaminophen 500 mg.....	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

#### Purposes

#### Uses

temporarily relieves these symptoms due to a cold:

- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- in a child under 12 years of age

- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

When using this product do not exceed recommended dosage

3124



**nighttime**  
severe cold  
and cough



**daytime**  
multi-symptom  
severe cold

**Drug Facts (continued)****Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 6 packets in 24 hours unless directed by a doctor.

**Drug Facts (continued)**

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

■ dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.

■ if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

**Other information**

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylketonurics: contains phenylalanine 22 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

**Questions? Call 1-888-547-7400**

**Nighttime Severe Cold And Cough****Drug Facts****Active ingredients (in each packet)****Purposes**

Acetaminophen 650 mg.....Pain reliever/fever reducer  
Diphenhydramine HCl 25 mg.....Antihistamine/cough suppressant  
Phenylephrine HCl 10 mg.....Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - minor sore throat pain
  - nasal and sinus congestion
  - headache
  - itchy nose or throat
  - runny nose
  - sneezing
  - itchy, watery eyes due to hay fever
  - cough due to minor throat and bronchial irritation
- temporarily reduces fever

**Warnings**

**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
  - blisters
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

- Do not use ■ in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ with any other product containing diphenhydramine, even one used on skin

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- Ask a doctor before use if you have
- high blood pressure
  - liver disease
  - diabetes
  - glaucoma
  - thyroid disease
  - heart disease
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema

**Drug Facts (continued)****Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- do not exceed recommended dosage
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose
adults and children 12 years of age and over	one packet
children under 12 years of age	do not use

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.

- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

**Other information**

- each packet contains: potassium 10 mg and sodium 25 mg
- phenylketonurics: contains phenylalanine 13 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, D&C yellow #10, FD&C blue #1, FD&C red #40, flavors, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

**Questions? Call 1-888-547-7400**

\*These products are not manufactured or distributed by GSK Consumer Healthcare, distributor of Theraflu® Severe Cold and Theraflu® Nighttime Severe Cold & Cough.

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

Do not use if printed packets are torn or punctured



Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in Mexico  
TM & ©2021 Target Brands, Inc.



[illegible]